

Assessment of Health-Related Quality of Life in Patients Receiving Regular Hemodialysis

Mahmoud A. Abdelghany^{1*}, Eman E. Elgohary² and Yasser A. Nienaa³

¹Nephrology specialist, Assistant Professor of Internal Medicine and Nephrology, Alexandria University, Egypt

²Professor of Internal Medicine and Nephrology, Alexandria University, Egypt

³Nephrology Department, Faculty of medicine, Alexandria University, Egypt

Abstract

This cross-sectional descriptive study was carried out at the hemodialysis unit of Karmouz Health Insurance Hospital, Alexandria, Egypt. The study was conducted on adult End-stage renal disease (ESRD) patients maintained on regular hemodialysis treatment for ≥ 1 year. Aim was to assess self-perception of ESRD patients undergoing regular Hemodialysis (HD) of their health-related quality of life (HRQOL) and explore resulted scores using kidney disease quality of life short form 36 instrument (KDQOL SF-36). A total of 81 eligible patients joined the study (49 males and 32 females). Aged 50.6 ± 12.7 years. Results showed that HRQOL of HD patients was very poor in all domains. The mean total score was below 50 (out of 100 point) with mean Physical Health Composite (PHC) = 35.57 ± 7.34 and mean Mental Health Composite (MHC) = 36.76 ± 10.22 . In general, scores were relatively higher in males, younger age groups, high educational level, employed and/or higher family income. Regarding clinical profile, this study showed that scores were lower among patients suffering anemia, dyslipidemia, hypoalbuminemia and/or those with comorbidities. Scores for general health survey were significantly lower in anemic patients (with hemoglobin level $< 11\text{g/dL}$). Working patients had significantly higher scores than non-working or retired. The study also showed that employment was the only significant predictor for total HRQOL score in a multiple linear regression model was designed to predict total score, while controlling for hemoglobin level, age, duration of being on HD ($p = 0.028$).

Keywords: End stage renal disease; Health related quality of life; Hemodialysis; Kidney disease quality of life short form; Anemia; Employment

Introduction

End-stage renal disease is a non-curable condition imposes a considerable burden on patients and their families [1,2]. It's been proven that the patient's perception is more important than the clinical assessment in determining HRQOL. Culture and values plays a vital role in shaping perception of HRQOL [3]. Although advances in HD treatment have contributed to improved patients' survival, they're still subjected to multiple threats and stressors that may worsen their HRQOL [4]. Lower HRQOL scores were strongly associated with higher morbidity and mortality [3]. The aim of the present study was to assess self-perception of patients with ESRD undergoing regular hemodialysis of HR-QOL, to study their epidemiological features, assess HR-QOL scores and explore its clinical correlates.

Patients and Methods

This cross-sectional descriptive study was carried out in September 2015 at the hemodialysis unit of Karmouz Health Insurance Hospital, Alexandria, Egypt. Among all adult ESRD patients aged > 18 years and maintained on regular hemodialysis treatment for ≥ 1 year, 81 eligible patients joined the study. (As exclusion criteria were: dementia, psychotic disorder, malignancy, active infection(s), acute medical or surgical condition, hospitalization within the prior month and physical or mental handicaps). A validated Arabic form of kidney disease quality of life short form (KD QOL SF-36) version 1.3 questionnaire was used as a self-report instrument to measure HR-QOL. Patients were encouraged to complete the questionnaire at the field of study and not at home, however some patients insisted to take it home and bring it back the next session. Most patients completed the questionnaire on their own, some patients -for various reasons e.g. being illiterate-completed the survey with assistance of the researcher or a coworker. The KDQOL Instrument is a self-report measure developed for individuals with kidney disease and on dialysis. It captures their

perceptions about concerns of symptoms or problems, effects and burden of kidney disease, work status, sleep, cognitive, social and sexual function. It also measures additional quality of life scales include: social support, dialysis staff encouragement and patient satisfaction [5]. The item related to sexual activity was considered it as an optional item to respect patients' privacy. It was missing in half of the patients yet we took that into consideration while doing analysis of the data.

All collected data checked for validity. Scoring was done to obtain universal scores ranging from 0 to 100, as higher scores indicate better QOL. Data entry and statistical analysis were performed using SPSS 18.0 statistical software package (SPSS Inc., Chicago, IL, USA). The distributions of quantitative variables were tested for normality using Kolmogorov-Smirnov test, which revealed that the data were normally distributed. Correlation between quantitative variables done using Pearson correlation test. Comparing quantitative variables between 2 groups of cases was conducted using independent t-test. Comparing quantitative variables between 3 groups or more of cases was conducted using oneway ANOVA. Kruskal Wallis test was used when sample size per group was small. Statistical significance was set at $P < 0.05$.

Results

Among the 81 HD patients studied (49 males + 32 females) aged 50.6 ± 12.7 years, HRQOL scores were poor in all domains and subscales. The mean total score was 47.2 ± 25.2 , mean Physical Health Composite (PHC) = 35.57 ± 7.34 and mean Mental Health Composite

***Corresponding author:** Mahmoud A. Abdelghany, Nephrology Specialist, Assistant Professor of Internal Medicine and Nephrology, Alexandria University, Egypt, Tel: 20 100 680 795; E-mail: amamh2@hotmail.com

Received: March 18, 2016; **Accepted:** April 02, 2016; **Published:** April 09, 2016

Citation: Abdelghany MA, Elgohary EE, Nienaa YA (2016) Assessment of Health-Related Quality of Life in Patients Receiving Regular Hemodialysis. J Nephrol Ther 6: 246. doi:[10.4172/2161-0959.1000246](https://doi.org/10.4172/2161-0959.1000246)

Copyright: © 2016 Abdelghany MA, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

(MHC) = 36.76 ± 10.22 . Regarding the ESRD- targeted areas, the mean total score was 54.48 ± 25.65 , the highest score was for dialysis staff encouragement scale (73.15 ± 23.64) followed by social support scale (72.22 ± 25.14) and quality of social interaction (71.52 ± 18.74). However, burden of kidney disease and work status scales yielded the lowest scores (25.62 ± 26.62 and 30.86 ± 37.56 respectively). Sleep scale score was 50.22 ± 21.27 . Only 40 patients responded to sexual activity inquiries and scored 66.88 ± 32.71 on sexual function scale (Table 1). On the 36-item health survey, the mean total score was 36.5 ± 25.67 . Emotional well-being and physical functioning scales yielded the highest score but were below 50 (48.3 ± 23.04 and 46.91 ± 27.79 respectively). The lowest scores represented the role limitations caused by physical and emotional health problems (21.6 ± 31.56 and 24.69 ± 35.27). Pain scale score was 40.31 ± 23.18 . Finally, the score overall health rating scale was 53.09 ± 16.63 . On comparing scores in both sexes (Table 1), females' scores were lower than males in mean total score (44.9 ± 23.7 and 49 ± 25.7 respectively) and all scales, except for the sexual function, social support scales and overall health rating item where female scores were relatively higher (Figures 1 and 2).

Unemployed and/or anemic patients had statistically significant reduced HRQOL ($P = 0.03$, $P = 0.02$ respectively) (Table 2). Scores varied according to age, sex, marital status, education level, presence of co-morbidities, dyslipidemia and hypoalbuminemia but without statistical significance. Multiple linear regression analysis for scores demonstrated that working status was the only significant predictor (employed patients' scores were higher than unemployed ones by 8.648 units, while controlling other predictors ($p = 0.028$) (Table 3).

Discussion

Long-term HD treatment often results in financial troubles, loss of

Domains and subscales	Mean	Median	SD
ESRD- targeted areas			
Symptom/problem list	49.97	52.08	23.08
Effects of kidney disease	51.04	50	24.49
Burden of kidney disease	25.62	18.75	26.62
Work status	30.86	0	37.56
Cognitive function	62.96	60	21.55
Quality of social interaction	71.52	73.33	18.74
Sexual function	66.88	75	32.71
Sleep	50.22	47.50	21.27
Social support	72.22	83.33	25.41
Dialysis staff encouragement	73.15	75	23.64
Patient satisfaction	44.86	50	27.09
The 36-item Health Survey			
Physical functioning	46.91	45	27.79
Role limitations-physical	21.60	0	31.56
Pain	40.31	35	23.18
General health	34.14	30	21.09
Emotional well-being	48.30	48	23.04
Role limitations-emotional	24.69	0	35.27
Social function	42.59	50	24.27
Energy/fatigue	33.46	30	19.18
Overall health	53.09	50	16.63
Total score	47.22	43.65	25.21
SF-12 Physical Health Composite	35.57	35.33	7.34
SF-12 Mental Health Composite	36.76	35.97	10.22

Table 1: KDQOL SF-36 scores for all studied patients.




Figure 1: HRQOL scores for ESRD- targeted areas.




Figure 2: The general health survey scores.

freedom, dependence on health care personnel and caregivers, negative impacts on marital status, family, and social activities. Thus, quality of life should be measured and monitored for better understanding of patients' condition [6]. This can also help to identify subgroups with relatively poor perceived health and guide interventions to improve their situations and avoid further serious consequences [4]. Over the years, consensus has been established that HRQOL is a multidimensional concept. As such, assessment of HRQOL generally covers physical, social, and psychological domains [4]. Several studies proved that HD patients suffer poor HRQOL, despite continuing advancements in the treatment [3,4,7-14]. Their HRQL is still much lower than general population [12,15]. Baseline measures of HRQOL were reduced in CKD patients in proportion to the severity grade of CKD [16]. Caregivers for ESRD patients may be affected as well, having a family member on maintenance dialysis therapy creates family burdens greater than those experienced by having a family member with several other chronic conditions [17]. Sexual function item left optional in our study for social issues and expressed by half of the patients. Its score was relatively high (66.8 ± 23). The Dialysis Outcomes and Practice Patterns Study (DOPPS) study in 2004 noticed variation in sexual function scores worldwide with mean scores 66.7, 63.3 and 60.5 for Europe, Japan and USA respectively with statically significant difference between Europe and USA [15]. In the present study, the mean of HRQOL scores were relatively higher in younger age groups in all domains, yet there was no statistically significant difference. The association of age with HRQOL is quite complex and illustrates the complexity of the QOL concept [4]. Some studies conducted in different countries also demonstrated

Work status*	Statistics	HR QOL Scores			
		ESRD- targeted areas	General health survey	Overall health item	Total score
Working (n= 23)	$\bar{x} \pm S.D.$	61.77 ± 17.65	41.42 ± 17.63	57.83 ± 15.06	53.31 ± 16.13
	Median	65.91	33.31	50	52.93
	Min-Max	15.57-97.73	24.19-97	40-100	22.21-97.55
	Std. Error	3.68	3.68	3.14	3.36
	95%CI♦	[54.1,69.4]	[33.79,49.0]	[51.3,64.3]	[46.3,60.29]
Not retired and not working (n=34)	$\bar{x} \pm S.D.$	51.46 ± 14.89	34.74 ± 17.75	55.59 ± 17.27	44.78 ± 14.92
	Median	52.19	32.36	60	43.91
	Min-Max	21.35-90.91	2.38-83.54	10-100	13.82-88.42
	Std. Error	2.55	3.04	2.96	2.56
	95%CI♦	[46.3,56.66]	[28.55,40.9]	[49.6,61.6]	[39.6,49.99]
Retired and not working (n=24)	$\bar{x} \pm S.D.$	49.13 ± 11.48	34.28 ± 15.22	45 ± 14.74	42.72 ± 11.44
	Median	50.09	33.56	50	42.97
	Min-Max	18.62-68.94	1.13-72.58	10-60	14.56-65.76
	Std. Error	2.34	3.11	3.01	2.34
	95%CI♦	[44.3,53.98]	[27.85,40.7]	[38.8, 51.2]	[37.9,47.55]
Test Statistics**					
	f	4.91	1.35	4.52	3.66
	df	2	2	2	2
	P value	0.01	0.266	0.014	0.03

*Working group included full or part time workers, not working and not retired group included housewives, patients who were searching for a job or who can't work due to their medical condition.

**Oneway ANOVA

♦ 95% Confidence Interval for mean [upper bound, lower bound]

Table 2: Comparing HR QOL scores between patients according to their work status using oneway ANOVA.

Model	Unstandardized Coefficients		Standardized Coefficients	t	Sig.
	B	Std. Error			
(Constant)	24.137	10.399		2.321	0.023
Hb ^a	8.841	4.658	0.206	1.898	0.062
Age	0.214	0.159	0.184	1.342	0.184
HD duration	0.215	0.48	0.051	0.448	0.655
Retired ^b	-6.030	4.402	-0.187	-1.370	0.175
Working ^b	8.648	3.858	0.265	2.242	0.028

Dependent: Mean total score
R²= 0.15, Model F=2.54, p=0.035

a: reference category is Anemic

b: reference category is Not working

Table 3: Multiple linear regression model to for HR QOL total score of hemodialysis patients.

that age was strongly inversely associated with the physical domain scores. As age increases in the elderly, physical function of the body decreases [2,4,14,18-22]. A recent Saudi showed that age was a negative predictor of HRQOL score [2]. Concerning gender, many studies of QOL in patients with ESRD have shown that women have lower scores on different QOL tests than men. This was observed in both pre-dialysis and dialysis patients. The reasons for poor QOL in women are not clear and seem to be related to psychological and social factors, rather than the illness itself [12,23,24]. This was consistent with the present study. It is of interest that Bayoumi et al 2014 showed reduced QOL scores in male patients and male sex was a negative predictor of QOL score [2]. Various factors such as geographic areas, ethnicity, sample size may explain this gender differences of QOL in HD patients [25]. Regarding work status, the current study revealed that HD patients' perception for work function and role of physical limitation scores were relatively

low (30.86 ± 37.6 and 21.6 ± 31.6 respectively). Only 28% of patients were employed (working full time or part time), the remaining were not working either due to being retired, in working-age and searching for a suitable job or can't work due to physical limitation. Whatever the cause the HRQOL scores were higher among the working group of patients in mean total score and ESRD targeted areas. Retired group ranked the least score. This difference was statistically significantly. And on applying a multiple Linear regression model to predict total scores, working group's scores were higher than non-working group by 8.648 units of total score while controlling for anemia state, (anemic/non-anemic), patient age, duration of being on hemodialysis treatment (p=0.028). The potential to increase the number of dialysis patients who are employed exists if the barriers to employment that have been identified can be removed [26]. Patients who are satisfied with hemodialysis treatment and accepting it have better working capacity [22]. Directing rehabilitation efforts toward the population of dialysis patients who identify themselves as willing and able to work seems a worthwhile strategy to improve their QOL [26]. Many sociodemographic and clinical factors can influence HRQOL of HD patients. Current study findings proved that future studies with more comprehensive and informative tools to focus on these variables and help us improve our knowledge about HRQOL determinants and in turn this will reflect on clinical practice to enhance their poor QOL status.

Conclusion

All domains of HRQOL were reduced in HD patients. Anemia and unemployment adversely affected HRQOL scores.

References

1. Soliman AR, Fathy A, Roshd D (2012) The growing burden of end-stage renal disease in Egypt. Ren Fail 34: 425-428.

2. Bayoumi M, Al Harbi A, Al Suwaida A, Al Ghonaim M, Al Wakeel J, et al. (2013) Predictors of quality of life in hemodialysis patients. *Saudi J Kidney Dis Transpl* 24: 254-259.
3. Joshi VD (2014) Quality of life in end stage renal disease patients. *World J Nephrol* 3: 308-316.
4. Mollaoglu M (2013) Quality of Life in Patients Undergoing Hemodialysis. In: Suzuki H (eds.) *Hemodialysis*, p: 823-843.
5. Hays RD, Kallich JD, Mapes DL, Coons SJ, Amin N, et al. (1997) *Kidney Disease Quality of Life Short Form (KDQOL-SF), version 1.3: A manual for use and scoring*. RAND, Washington D.C.
6. Thaweethamcharoen T, Srimongkol W, Noparatayaporn P, Jariyayothin P, Sukthinthai N, et al. (2013) Validity and Reliability of KDQOL-36 in Thai Kidney Disease Patient. *Value in Health* 2: 98-102.
7. Abraham S, Venu A, Ramachandran A, Chandran PM, Raman S (2012) Assessment of quality of life in patients on hemodialysis and the impact of counseling. *Saudi J Kidney Dis Transpl* 23: 953-957.
8. Sathvik BS, Parthasarathi G, Narahari MG, Gurudev KC (2008) An assessment of the quality of life in hemodialysis patients using the WHOQOL-BREF questionnaire. *Indian J Nephrol* 18: 141-149.
9. Abdel-Kader K, Unruh ML, Weisbord SD (2009) Symptom burden, depression, and quality of life in chronic and end-stage kidney disease. *Clin J Am Soc Nephrol* 4: 1057-1064.
10. Jaar BG, Chang A, Plantinga L (2013) Can we improve quality of life of patients on dialysis? *Clin J Am Soc Nephrol* 8: 1-4.
11. Sesso R, Rodrigues-Neto JF, Ferraz MB (2003) Impact of socioeconomic status on the quality of life of ESRD patients. *Am J Kidney Dis* 41: 186-195.
12. Valderrábano F, Jofre R, López-Gómez JM (2001) Quality of life in end-stage renal disease patients. *Am J Kidney Dis* 38: 443-464.
13. Rambod M, Shabani M, Shokrpour N, Rafii F, Mohammadalliha J (2011) Quality of life of hemodialysis and renal transplantation patients. *Health Care Manag (Frederick)* 30: 23-28.
14. Stojanovic M, Stefanovic V (2007) Assessment of health-related quality of life in patients treated with hemodialysis in Serbia: Influence of comorbidity, age, and income. *Artif Organs* 31: 53-60.
15. Mapes DL, Bragg-Gresham JL, Bommer J, Fukuhara S, McKeivitt P, et al. (2004) Health-related quality of life in the Dialysis Outcomes and Practice Patterns Study (DOPPS). *Am J Kidney Dis* 44: 54-60.
16. Mujais SK, Story K, Brouillette J, Takano T, Soroka S, et al. (2009) Health-related quality of life in CKD Patients: correlates and evolution over time. *Clin J Am Soc Nephrol* 4: 1293-1301.
17. Belasco AG, Sesso R (2002) Burden and quality of life of caregivers for hemodialysis patients. *Am J Kidney Dis* 39: 805-812.
18. Blake C, Codd MB, Cassidy A, O'Meara YM (2000) Physical function, employment and quality of life in end-stage renal disease. *J Nephrol* 13: 142-149.
19. Simmons RG, Abress L (1990) Quality-of-life issues for end-stage renal disease patients. *Am J Kidney Dis* 15: 201-208.
20. Braga SF, Peixoto SV, Gomes IC, Acúrcio Fde A, Andrade EI, et al. (2011) Factors associated with health-related quality of life in elderly patients on hemodialysis. *Rev Saude Publica* 45: 1127-1136.
21. Moura A, Madureira J, Alija P, Fernandes JC, Oliveira JG, et al. (2015) Effect of Aging in the Perception of Health-Related Quality of Life in End-Stage Renal Disease Patients under Online-Hemodiafiltration. *Aging Dis* 6: 17-26.
22. Anees M, Malik MR, Abbasi T, Nasir Z, Hussain Y, et al. (2014) Demographic factors affecting quality of life of hemodialysis patients - Lahore, Pakistan. *Pak J Med Sci* 30: 1123-1127.
23. Mingardi G, Cornalba L, Cortinovis E, Ruggiata R, Mosconi P, et al. (1999) Health-related quality of life in dialysis patients. A report from an Italian study using the SF-36 Health Survey. *DIA-QOL Group. Nephrol Dial Transplant* 14: 1503-1510.
24. Moreno F, Gomez JML, Jofre R, Valderrabano F, Gonzalez L, et al. (1996) Quality of life in dialysis patients. A Spanish multicentre study *Nephrol Dial Transplant* 11: 125-129.
25. Einollahi B, Motalebi M (2014) Remarks about the study of predictors of quality of life in hemodialysis patients. *Saudi J Kidney Dis Transpl* 25: 881-882.
26. Curtin RB, Oberley ET, Sacksteder P, Friedman A (1996) Differences between employed and nonemployed dialysis patients. *Am J Kidney Dis* 27: 533-540.

Citation: Abdelghany MA, Elgohary EE, Nienaa YA (2016) Assessment of Health-Related Quality of Life in Patients Receiving Regular Hemodialysis. *J Nephrol Ther* 6: 246. doi:[10.4172/2161-0959.1000246](https://doi.org/10.4172/2161-0959.1000246)

OMICS International: Publication Benefits & Features

Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

Special features:

- 700 Open Access Journals
- 50,000 editorial team
- Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, EBSCO, Index Copernicus and Google Scholar etc
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: www.omicsonline.org/submit/